

BRYOPHYTE CHECKLIST AND EUROPEAN RED LIST OF THE BRUSSELS-CAPITAL REGION, FLANDERS AND WALLONIA (BELGIUM)

André SOTIAUX^{1,2,*}, Herman STIEPERAERE¹ and Alain VANDERPOORTEN²

¹ National Botanic Garden of Belgium, Domein van Bouchout, B - 1860 Meise, Belgium

² Université de Liège, Institut de Botanique, B22 Sart Tilman, B - 4000 Liège, Belgium

(* Author for correspondence)

Received 10 August 2006; accepted 26 October 2006.

ABSTRACT. — An amended checklist of the bryophytes of Belgium is presented. Although some taxonomic positions adopted in the most recent European checklist of mosses are discussed, the latter is used as a reference in an attempt to homogenize taxonomic concepts at a wide scale and enable comparisons across countries. Since the previous 2001 checklist of Belgian bryophytes, a total of 17 and 14 species must be added to the Belgian bryoflora owing to taxonomic changes and floristic discoveries, respectively. The report of *Grimmia caespiticia*, at an altitude of 500-600 m, is particularly noteworthy because this alpine species usually occurs above 2 000 m. Five additional taxa, *Scapania cuspiduligera*, *Dalytrichia fragilifolia*, *Ephemerum cohaerens*, *Orthotrichum shawii* and *Tortella inclinata* var. *densa*, are reported here as new to Belgium. Six species were previously erroneously reported from Belgium and are excluded. As a consequence, the bryophyte flora of Belgium consists of 748 species, including 5 hornworts, 173 liverworts, and 570 mosses whose occurrence in the three regions of Belgium, namely Flanders (5 hornworts, 122 liverworts, 399 mosses), Wallonia (3 hornworts, 169 liverworts, 546 mosses), and Brussels-Capital Region (46 liverworts, 196 mosses), is documented. Thirty-six species (6 liverworts and 30 mosses) have not been recorded since 1980 and are most probably extinct. The new checklist is used to update the list of European red list species that are present in Belgium, which now includes 49 species.

KEY WORDS. — Belgium, Bryophytes, checklist, red list species.

INTRODUCTION

The most recent publication of the European checklist of mosses (HILL *et al.* 2006), which incorporates substantial taxonomic changes largely based upon recent outcomes of the molecular systematics, will undoubtedly influence the taxonomic positions so far adopted in the different countries. As Hill's *et al.* work resulted from the joint effort of a panel of taxonomic experts, reaching a consensus at a large scale remains an extremely challenging task. A good example is offered by the conflictual

treatments of difficult families such as the Pottiaceae (contrast, e.g., ZANDER 1993, GUERRA & CANO 2000, and WERNER *et al.* 2005a,b). Another issue is that the status of some of the species retained in the European checklist has long been controversial because of the strong overlap in morphological features and the existence of a range of puzzling intermediates. Examples of such cases are numerous (e.g., *Aloina aloides* and *A. ambigua*; *Plagiothecium succulentum* and *P. nemorale*; *P. curvifolium* and *P. laetum*; *Didymodon vinealis* and *D. insulanus*; *Hygroamblystegium tenax* and *H.*

varium; *Tortella bambgereri* and *T. tortuosa*; *Orthotrichum schimperi* and *O. pumilum*). While the genetic bases for justifying species recognition is lacking in most cases, it is clear that some of these species form complexes of polyphyletic origin (SHAW & ALLEN 2000, VANDERPOORTEN *et al.* 2004) and convergent morphology (VANDERPOORTEN & JACQUEMART 2004). In such cases, the conflicting interpretations of taxonomists will still lead to substantial differences in diversity assessments and hence, difficulties of comparisons across- (and even within-) countries. Thus, although Hill's *et al.* reference list should arguably serve as a reference basis to homogenize taxonomic concepts between countries, departures from the proposed list will still persist in the checklist adopted within each country. In The Netherlands for example (SIEBEL *et al.* 2005), most of the above species have been synonymized.

In this paper, we discuss the application of the new European checklist to the Belgian bryoflora. While preparing this revision, it also appeared that, to be useful to a broad panel of users including local florists, taxonomists seeking comparative data at a national or regional scale, and conservation agencies, the checklist should not only provide a taxonomic and nomenclatural reference basis for the Belgian bryoflora, but also include information about species occurrence at the regional level and threats. Our motivation for presenting an updated checklist since our previous contribution (SOTIAUX & VANDERPOORTEN 2001) was further stimulated by a number of additional species reports and the important con-

sequences for the European red list species present in Belgium (ECCB 1995).

METHODS

For liverworts and hornworts, nomenclature follows GROLLE & LONG (2000) with some modifications due to subsequent taxonomic and nomenclatural works, including GROLLE & SO (2003) for *Metzgeria* and SZWEJKOWSKI *et al.* (2005) for *Conocephalum*. Following DE ROO *et al.* (2007), ENGEL & BRAGGINS (2005) are not followed in accommodating *Mylia anomala* within its own genus *Leiomylia*. GROLLE & LONG (2000), who treat *Lophocolea* and *Chiloscyphus* as separate genera, are also followed. There is, however, growing evidence from molecular studies (e.g., HENTSCHEL *et al.* 2006) that ENGEL & SCHUSTER (1984) were right in merging *Lophocolea* with *Chiloscyphus*.

For mosses, the European checklist of HILL *et al.* (2006) served as a reference. Therefore, HILL *et al.* (2006) are followed in merging a series of species (e.g., *Bryum barnesii* and *B. dichotomum*), even though they display, at least in Belgium, a clear morphological discontinuity and might deserve a distinct taxonomic status. For the same reasons, HILL *et al.* (2006) are followed in recognizing a series of morphologically extremely similar and intergrading species (see above), although we believe that such species, because of the difficulty in their identification, add 'noise' to floristic inventories and hence, that a more conservative strategy would be to reduce these species to synonymy. In the case of *Leucobryum*, however, HILL *et al.* (2006) are not followed in recognizing three species in the European flora for reasons detailed in the text. Nomenclatural changes since the 2001 checklist are summarized in Table 1.

Table 1. List of nomenclatural changes, with corresponding synonymies, since the previous check-list of the bryophytes of Belgium by SOTIAUX & VANDERPOORTEN (2001).

Nomenclature of SOTIAUX & VANDERPOORTEN (2001)	Current name
Liverworts	
<i>Leiocolea bantriensis</i>	<i>Leiocolea collaris</i>
<i>Metzgeria fruticulosa</i>	<i>Metzgeria violacea</i>
Mosses	
<i>Amblystegium fluviatile</i>	<i>Hygroamblystegium fluviatile</i>
<i>Amblystegium humile</i>	<i>Hygroamblystegium humile</i>
<i>Amblystegium tenax</i>	<i>Hygroamblystegium tenax</i>
<i>Amblystegium varium</i>	<i>Hygroamblystegium varium</i>
<i>Aphanorhynchum patens</i>	<i>Physcomitrella patens</i>
<i>Brachythecium oedipodium</i>	<i>Sciuro-hypnum oedipodium</i>

<i>Brachythecium oxycladum</i>	<i>Brachythecium laetum</i>
<i>Brachythecium plumosum</i>	<i>Sciuro-hypnum plumosum</i>
<i>Brachythecium populeum</i>	<i>Sciuro-hypnum populeum</i>
<i>Brachythecium reflexum</i>	<i>Sciuro-hypnum reflexum</i>
<i>Brachythecium velutinum</i>	<i>Brachytheciastrum velutinum</i>
<i>Bryum barnesii</i>	<i>Bryum dichotomum</i>
<i>Bryum imbricatum</i>	<i>Bryum archangelicum</i>
<i>Bryum rutilans</i>	<i>Bryum pallens</i>
<i>Bryum stirtonii</i>	<i>Bryum elegans</i>
<i>Bryum subelegans</i>	<i>Bryum moravicum</i>
<i>Bryum versicolor</i>	<i>Bryum dichotomum</i>
<i>Calliergon sarmenosum</i>	<i>Warnstorfia sarmentosa</i>
<i>Cirriphyllum tommasinii</i>	<i>Brachythecium tommasinii</i>
<i>Cynodontium polycarpum</i>	<i>Cynodontium polycarpon</i>
<i>Dicranella palustris</i>	<i>Dichodontium palustre</i>
<i>Dicranum bergeri</i>	<i>Dicranum undulatum</i>
<i>Ditrichum cylindricum</i>	<i>Trichodon cylindricus</i>
<i>Eurhynchium crassinervium</i>	<i>Cirriphyllum crassinervium</i>
<i>Eurhynchium flotowianum</i>	<i>Sciuro-hypnum flotowianum</i>
<i>Eurhynchium hians</i>	<i>Oxyrrhynchium hians</i>
<i>Eurhynchium praelongum</i>	<i>Kindbergia praelonga</i>
<i>Eurhynchium pulchellum</i>	<i>Eurhynchiastrum pulchellum</i>
<i>Eurhynchium pumilum</i>	<i>Oxyrrhynchium pumilum</i>
<i>Eurhynchium schleicheri</i>	<i>Oxyrrhynchium schleicheri</i>
<i>Eurhynchium speciosum</i>	<i>Oxyrrhynchium speciosum</i>
<i>Eurhynchium striatum</i>	<i>Plasteurhynchium striatum</i>
<i>Fissidens incurvus</i>	<i>Fissidens viridulus</i> var. <i>incurvus</i>
<i>Funaria muhlenbergii</i>	<i>Entosthodon muhlenbergii</i>
<i>Funaria pulchella</i>	<i>Entosthodon pulchellus</i>
<i>Hylocomium brevirostre</i>	<i>Loeskeobryum brevirostre</i>
<i>Hylocomium pyrenaicum</i>	<i>Hylocomiastrum pyrenaicum</i>
<i>Hylocomium umbratum</i>	<i>Hylocomiastrum umbratum</i>
<i>Hypnum lacunosum</i>	<i>Hypnum cupressiforme</i> var. <i>lacunosum</i>
<i>Hypnum pratense</i>	<i>Breidleria pratensis</i>
<i>Hypnum resupinatum</i>	<i>Hypnum cypresiforme</i> var. <i>resupinatum</i>
<i>Leucobryum juniperoides</i>	<i>Leucobryum glaucum</i>
<i>Microbryum curvicolle</i>	<i>Microbryum curvicollum</i>
<i>Octodiceras fontanum</i>	<i>Fissidens fontanus</i>
<i>Pleuridium palustre</i>	<i>Cleistocarpidium palustre</i>
<i>Polytrichum formosum</i>	<i>Polytrichastrum formosum</i>
<i>Polytrichum longisetum</i>	<i>Polytrichastrum longisetum</i>
<i>Pseude-calliergon lycopodioides</i>	<i>Pseudocalliergon lycopodioides</i>
<i>Schistidium singarens</i>	<i>Schistidium helveticum</i>
<i>Scleropodium purum</i>	<i>Pseudoscleropodium purum</i>
<i>Seligeria paucifolia</i>	<i>Seligeria calycina</i>
<i>Sematophyllum micans</i>	<i>Hageniella micans</i>
<i>Serpoleskea confervoides</i>	<i>Amblystegium confervoides</i>
<i>Sphagnum denticulatum</i>	<i>Sphagnum auriculatum</i>
<i>Syntrichia inermis</i>	<i>Tortula inermis</i>
<i>Syntrichia intermedia</i>	<i>Syntrichia montana</i>
<i>Syntrichia pagorum</i>	<i>S. laevipila</i>
<i>Syntrichia ruraliformis</i>	<i>Syntrichia ruralis</i> var. <i>ruraliformis</i>
<i>Thuidium abietinum</i>	<i>Abietinella abietina</i>
<i>Thuidium philibertii</i>	<i>Thuidium assimile</i>
<i>Tortula acaulon</i>	<i>Phascum cuspidatum</i>
<i>Tortula protobryoides</i>	<i>Protobryum bryooides</i>
<i>Trichostomopsis australasiae</i>	<i>Didymodon australasiae</i>

RESULTS

CHANGES WITH THE PREVIOUS CHECKLIST

New taxa

Fourteen species are reported as new since the 2001 checklist: *Haplomitrium hookeri* (SOTIAUX *et al.* 2006), *Scapania calcicola* (SOTIAUX *et al.* 2002), *S. cuspiduligera* (see below), *Bryum dyfrynnense* (DE BEER & MARTENS 2007), *Dalytrichia fragilifolia* (see below), *Ephememerum cohaerens* (see below), *Grimmia caespiticia* (MUÑOZ & PANDO 2000), *G. dissimulata* (DE BEER 2007), *G. elongata* (GREVEN & SOTIAUX 2003), *Isothercium holtii* (cited in SIEBEL & DURING 2006 for the Ardennes district, based on Siebel 01.381, Robertville, Warche Valley, IFBL G8.34.23, 7.08.2001, BR and herb. Siebel), *Orthotrichum shawii* (see below), *Polytrichastrum alpinum* (REINEKE *et al.* 2002), *Rhytidadelphus subpinnatus* (SCHUMACKER & SOTIAUX 2003) and *Tortula mucronifolia* (SOTIAUX & SOTIAUX 2005).

Grimmia caespiticia (Brid.) Jur. was once collected in Belgium at the beginning of the 20th century (Cornet, 1906, NY!) (MUÑOZ & PANDO 2000). The collection includes two fertile plants with exserted capsules and an obliquely rostrate operculum. The leaves are distinctly longitudinally furrowed. Hence, the specimen perfectly matches the circumscription of *G. caespiticia*. The Belgian locality, at an altitude of 500-600 m, is completely outside the altitudinal range of this alpine species, which mostly occurs at an altitude of 2 000 m or higher, with a few occurrences reported down to an altitude of 1 400 m (GREVEN 1995).

Scapania cuspiduligera, *Dalytrichia fragilifolia*, *Ephememerum cohaerens*, *Orthotrichum shawii* and *Tortella inclinata* var. *densa* are reported here as new to Belgium.

Scapania cuspiduligera: Prov. Namur, Durval, Bocq valley (IFBL H5.28.23), on rocks in old quarry together with *Encalypta streptocarpa* and *Rhynchostegium murale*, alt. 175 m, 28.10.1979 (herb. Sotiaux, Sotiaux 3).

Dalytrichia fragilifolia: Prov. Namur, Dinant, Meuse valley towards Colébi (IFBL H5.57.12), shaded calcareous rocks, alt. 100 m, 08.06.1986 (herb. Sotiaux, Sotiaux 5450).

Ephememerum cohaerens: Prov. Liège, Herrenrath (IFBL F8.13.21), wet sandy track in an ancient sand quarry, with *Calliergonella cuspidata*, alt. 260 m, 01.10.2006 (LG, Vanderpoorten B1362); Prov. Namur, Merlemon (IFBL J5.12.13), wet calcareous ground of an ancient limestone quarry with *Pellia endiviifolia*, alt. 250 m, 27.11.2001 (herb. Sotiaux, Sotiaux 27619, herb. J.-P. Duvivier, Duvivier s.n.).

Orthotrichum shawii: Prov. Liège, Büllingen, Our valley between Hüllscheid and Berterath (IFBL H9.11.23), epiphytic on *Fraxinus excelsior* at 520 m, 25.06.2005 (herb. Sotiaux, Sotiaux 32729).

Tortella densa: Prov. Liège, Huy (IFBL G6.14.43), calcareous gravelly ground, together with *Campyliadelphus chrysophyllus*, *Didymodon fallax*, and *T. inclinata* var. *inclinata*, alt. 85 m, 10.11.2005 (LG, Vanderpoorten B899 and herb. Sotiaux, Sotiaux 33392). As opposed to *T. inclinata*, the dorsal side of the costa in *T. densa* is not covered by a layer of isodiametric cells. Although this is a key character in the taxonomy of the genus (e.g., SMITH 2004), *T. densa* is now treated as a variety of *T. inclinata* in HILL *et al.* (2006). This taxon is therefore not included in the checklist, wherein we do not list infraspecific taxa.

A total of 17 species must be added to the checklist due to the creation or erection at the species level of former infraspecific taxa, already known from Belgium. These species include *Conocephalum salebrosum*, *Aloina ambigua*, *Anomobryum concinnum*, *Bryum kunzei*, *Campylium protensum*, *Dichodontium flavescens*, *Didymodon insulanus*, *Ephememerum minutissimum*, *Heterocladium flaccidum*, *Orthotrichum schimperi*, *Palustriella falcata*, *Plagiothecium succulentum*, *Polytrichum uliginosum*, *Syntrichia calcicola*, *Tortella bambergeri*, *Tortula schimperi*, and *Zygodon stirtonii*.

Rediscovered taxa

Two species considered extinct have been rediscovered.

Philonotis marchica had not been seen in Belgium since 1912, (RAEYMAEKERS 1981) but

was discovered in a wetland restoration area near Bruges (STIEPERAERE 2007).

Pterygoneurum lamellatum was found in 2005 very close to the Dutch and German border in Bassenge (ANDRIESSEN *et al.* 2006). The previous collection of the species dated back to more than 100 years ago.

Excluded taxa

Three species must be deleted from the previous checklist as a result of the reduction to synonymy in HILL *et al.* (2006). These species include *Bryum barnesii* (=*B. dichotomum*), *B. rutilans* (=*B. pallens*), and *B. versicolor* (=*B. dichotomum*). A fourth species, *Leucobryum juniperoides*, is also deleted. Contra FRAHM (2005), which is the reference cited by HILL *et al.* (2006) for keeping *L. juniperoides*, *L. albidum* is by no means characterized by leaves that are less than 2 mm long (2-4 mm for example reported by CRUM & ANDERSON 1981). *Leucobryum albidum* does intergrade morphologically with *L. glaucum*, the latter including in its range the poorly defined *L. juniperoides*. *Leucobryum juniperoides* was therefore synonymized and it was proposed that the European *Leucobryum* flora consists of only two species, *L. glaucum* and *L. albidum* (VANDERPOORTEN *et al.* 2003). Only the former has to date been found in Belgium. Although *L. albidum* exhibits a more southwestern distribution than *L. glaucum*, it was found in France and Germany, so that the species is very likely to occur in Belgium as well.

Seven additional species are excluded for the reasons detailed below.

Scapania praetervisa. The occurrence of this boreal, calcicolous species on highly acidic soils with *Calluna vulgaris* and other strictly acidophilous species such as *Lophozia grandiretis* (DE ZUTTERE & POHL 1991) at low elevation is ecologically unlikely. The presence of pinkish gemmae, used as evidence for the identity of the collection with *S. praetervisa* (DE ZUTTERE & POHL 1991), can be observed in *S. scandica*, an acidophilous species present in the Ardennes. The material of *S. praetervisa* is unfortunately unavailable for re-examination and the species is,

therefore, tentatively excluded from the Belgian checklist.

Fissidens curnovii (=*F. bryoides* var. *caespitans*). All the specimens attributed to this taxon from Belgium were re-examined by M. A. Bruggeman and proved to belong to *F. bryoides* var. *bryoides*.

Fissidens limbatus (=*F. crispus*). No collection of this Mediterranean species was found at BR or LG and the record most probably resulted from a synonymy error in SCHUMACKER *et al.* (1985).

Grimmia reflexidens Müll. Hal. was reported from Belgium from the same locality as *G. caespiticia* in Spa, Promenade Annette et Lubin (MUÑOZ & PANDO 2000). The collection of *G. reflexidens* (Cornet, 1906, BP!) is sterile and the gametophyte does not match that of the typical expression of the species. MUÑOZ & PANDO (2000) indicated that the group of species including *G. australis* (Dixon & Sainsbury) J. Muñoz & Ochyra, *G. alpestris* (F. Weber & D. Mohr) Schleicher., and *G. reflexidens* is characterized by bulging laminal cells in leaf cross-section, but emphasized that this character cannot be observed on poor specimens. Pending convincing evidence of the presence of *G. reflexidens* in Belgium based on fertile material, we exclude this alpine species from the checklist.

Grimmia mollis Bruch & Schimp., reported from Belgium (MUÑOZ & PANDO 2000), was tentatively not included in the previous checklist of bryophytes of Belgium pending further evidence of the presence of this alpine species in Belgium (SOTIAUX & VANDERPOORTEN 2001). J. Muñoz (pers. comm.) confirmed that the report of *G. mollis* in Belgium was a mistake.

Meesia longisetata. DE SLOOVER (1970) renamed the original collection as *Aulacomnium palustre*. A later collection at BR from Troch's herbarium is of doubtful origin and probably a fraud.

Meesia uliginosa. KICKX (1867) reported fragments of this species from a supply of *Sphagnum* for the Botanic Garden of Ghent harvested near Damme in 1857. DE SLOOVER (1970) showed that the specimen in BR with a label in Troch's handwriting (Damme, Kickx, 1850) is probably a fraud. We found that the collection is

mixed with *Blepharostoma trichophyllum* and has mica fragments in the peaty substrate. This makes it extremely unlikely that the specimen comes from anywhere near Damme.

Taxa considered as extinct

Catoscopium nigritum. This species was treated as extinct in our previous checklist. Evidence for its presence in Belgium remains, however, subject to controversy. There has been no record of this species from Flanders since DE SLOOVER (1970) re-identified the original specimen cited in DELOGNE (1883) as *Dicranella varia* and showed that a second, correctly named specimen in BR from the Troch collection is most probably a fraud. In the herbarium of the Carnoy Institute (LV, transferred to BR in 1999), we found another specimen of *C. nigritum*. The species name and the collector's name are in Dellogne's handwriting; the locality "Entre Ostende & Blanckenbergh" is a later addition in a different handwriting. Therefore, some doubts remain regarding this record. Most recently, a collection made by Castagne in 1956 from Baraque Michel in the Ardennes was found in the Parent herbarium (DE ZUTTERE & SOTIAUX 2007). In the complete absence of basic fens around Baraque Michel, however, a little doubt also persists regarding the validity of this record.

THE CHECKLIST

As a consequence of the changes exposed above, the bryophyte flora of Belgium (30,528 km²) consists of 748 species, including 5 hornworts, 173 liverworts, and 570 mosses (Appendix). A total of 526 species is recorded to date from Flanders (13,522 km²), including 5 hornworts, 122 liverworts, and 399 mosses. The Brussels-Capital Region (162 km²) bryophyte flora is composed of 242 species, including 46 liverworts and 196 mosses. In Wallonia (16,844 km²), 718 species have been recorded to date, including 3 hornworts, 169 liverworts, and 546 mosses (Appendix).

Thirty-six species, including 6 liverworts and 30 mosses, have not been recorded since 1980 and are probably extinct.

CONSEQUENCES FOR THE EUROPEAN RED LIST SPECIES OF BELGIUM

Thirty-six European red-list species were listed as occurring in Belgium (ECCB 1995) [excluding *Didymodon glaucus*, only reported from the Grand-Duchy of Luxembourg, which is subject to a separate survey (WERNER 2003)]. Four species: *Fossombronia incurva*, *Grimmia lisae*, *Hamatocaulis vernicosus* and *Rhynchosstegium rotundifolium*, which do occur in Belgium, were omitted from the 1995 ECCB list. Since the publication of the European red list, twelve additional red list species have been reported from Belgium: *Anthoceros caucasicus* (HEYLEN *et al.* 2001), *Haplomitrium hookeri* (SOTIAUX *et al.* 2006), *Ephemerum stellatum* (VANDERPOORTEN *et al.* 1996), *Orthotrichum consimile* (SOTIAUX *et al.* 1998a), *O. ibericum* (HEYLEN & WALRAEVENS 2001), *O. rogeri* (SOTIAUX & SOTIAUX 2002), *O. scanicum* (SOTIAUX *et al.* 1998b), *Schistidium helveticum* (BLOM 1996), *S. papillosum* (BLOM 1996), *S. pruinorum* (BLOM 1996), and *Ulota rehmannii* (SOTIAUX & SOTIAUX 1999). Two additional species, *Ephemerum cohaerens* and *Grimmia caespiticia*, are reported here. Furthermore, *Aneura maxima* was found new to Europe in Belgium (ANDRIESEN *et al.* 1995). We have since observed the species fairly frequently in the Belgian and French parts of the Ardennes, where it grows in deep seeps, often with *Chrysosplenium*. It seems to be an overlooked species that may have been confounded with *Pellia neesiana* and *P. epiphylla*. At the European scale, however, *Aneura maxima* is only reported from 1-2 localities in France, Luxembourg, Poland, and Finland (VANDERPOORTEN *et al.* 2006). As a consequence, we suggest with WERNER (2003) that the species should be included within the European red list, but with the status 'insufficiently known', as its exact distribution in Europe remains to be documented (VANDERPOORTEN *et al.* 2006). Lastly, five species must be deleted from the European red list species of Belgium. *Buxbaumia viridis* (DE SLOOVER & STIEPERAERE 1999) and *Fissidens ovatifolius* were erroneously listed from Belgium in the treatment of ECCB (1995). *Meesia longiseta* must be excluded from the checklist of

Belgium for reasons detailed above, while *Bryum stirtonii*, *B. versicolor*, and *Rhynchostegiella jacquinii* are currently considered to be conspecific with *Bryum elegans*, *B. dichotomum*, and *R. teneriffae*, respectively.

As a consequence, 49 species listed in the European red list (ECCB 1995) occur in Belgium (Table 2). Seven species are probably extinct in

Belgium and all the others are known from one to few localities, except *Aneura maxima* (VANDERPOORTEN *et al.* 2006), *Cleistocarpidium palustre*, and *Orthotrichum consimile* (SOTIAUX *et al.* 1998a, SOTIAUX & VANDERPOORTEN 2004), for which the Belgian populations play a significant role in the conservation of those species in Europe.

Table 2. The European red list species (ECCB 1995) present in Belgium.

Species	IUCN category	Status in Belgium
<i>Anthoceros caucasicus</i>	R	Two localities (STIEPERAERE 2002)
<i>Aneura maxima</i>	K	Locally fairly common (VANDERPOORTEN <i>et al.</i> 2006)
<i>Cephaloziella elachista</i>	K	
<i>Fossumbronia incurva</i>	R	
<i>Haplomitrium hookeri</i>	R	A few shoots found in a single locality (SOTIAUX <i>et al.</i> 2006)
<i>Pallavicinia lyellii</i>	V	
<i>Riccia huebeneriana</i>	R	
<i>Amblystegium radicans</i>	R	
<i>Anomodon rostratus</i>	R	One locality (SCHUMACKER <i>et al.</i> 1982)
<i>Brachythecium laetum</i>	R	
<i>Bryum neodamense</i>	R	Not seen since the 19th century
<i>Bryum tenuisetum</i>	K	
<i>Bryum warneum</i>	R	
<i>Campylostelium saxicola</i>	R	Not seen for >100 years (SOTIAUX & VANDERPOORTEN 2004)
<i>Cleistocarpidium palustre</i>	R	Known from >50 localities, probably overlooked due to its short, seasonal presence
<i>Dicranum viride</i>	V	Not seen since 1980
<i>Ephemerum cohaerens</i>	E	Reported from two localities, including one found in 2006; apparently exceedingly rare but possibly somewhat overlooked
<i>Ephemerum recurvifolium</i>	R	
<i>Ephemerum sessile</i>	R	
<i>Ephemerum stellatum</i>	V	Disappeared because of the spreading of dolomite on its only locality (VANDERPOORTEN <i>et al.</i> 2001)
<i>Fissidens arnoldii</i>	R	
<i>Fissidens monguilloni</i>	R	Reported from many localities (DE ZUTTERE 1993) based on sterile material but undistinguishable from the common <i>F. bryoides</i> in this state and probably truly very rare
<i>Grimmia atrata</i>	R	
<i>Grimmia caespiticia</i>	R	One locality, not seen for >100 years
<i>Grimmia lisae</i>	R	
<i>Hamatocaulis vernicosus</i>	K	
<i>Leptodontium gemmascens</i>	R	Very rare and decreasing (DE ZUTTERE <i>et al.</i> 2002)
<i>Microbryum floerkeanum</i>	K	
<i>Micromitrium tenerum</i>	V	
<i>Mielichhofferia mielichhofferiana</i>	K	
<i>Orthotrichum consimile</i>	Ev	>20 localities
<i>Orthotrichum ibericum</i>	K	

<i>Orthotrichum rogeri</i>	V	
<i>Orthotrichum scanicum</i>	E	Still very rare but increasingly found
<i>Orthotrichum sprucei</i>	R	
<i>Physcomitrium sphaericum</i>	R	
<i>Pterygoneuron lamellatum</i>	V	
<i>Rhynchostegiella tenuicaulis</i>	K	
<i>Rhynchostegium rotundifolium</i>	R	
<i>Schistidium helveticum</i>	K	
<i>Schistidium papillosum</i>	K	
<i>Schistidium pruinosum</i>	K	
<i>Schistidium trichodon</i>	K	
<i>Sematophyllum demissum</i>	R	Not seen since 1978
<i>Tortula cernua</i>	R	One 19 th century locality on a carbide dump
<i>Ulota rehmanni</i>	E	A few specimens from a single locality (SOTIAUX & SOTIAUX 1999)
<i>Weissia rostellata</i>	R	
<i>Weissia squarrosa</i>	R	
<i>Zygodon forsteri</i>	V	Not seen for >100 years (SOTIAUX & VANDERPOORTEN 2004)

ACKNOWLEDGEMENTS

We are grateful to Henk Siebel for his constructive remarks on the manuscript. Many thanks are due to Beata Papp and Bill Buck for the loan of specimens and Jezus Muñoz for discussing the problems related to the occurrence of some *Grimmia* species in Belgium.

REFERENCES

- ANDRIESSEN L., SOTIAUX A., NAGELS C. & SOTIAUX O., 1995. — *Aneura maxima* (Schiffn.) Steph. in Belgium, new for the European liverwort flora. *J. Bryol.* **18**: 803-806.
- ANDRIESSEN L., HEUSERR M., NAGELS C. & VANDEKERCKHOVE K., 2006. — [http://users.skynet.be/natuur_in_limburg/
Bryolim/Verslagen/PLAM/Pterygoneurum.htm](http://users.skynet.be/natuur_in_limburg/Bryolim/Verslagen/PLAM/Pterygoneurum.htm)
- BLOM H.H., 1996. — A revision of the *Schistidium apocarpum* complex in Norway and Sweden. *Bryophytorum Bibliotheca* **49**: 1-333.
- CRUM H. & ANDERSON L.E., 1981. — *Mosses of eastern North America*. Columbia University Press, New York.
- DE BEER D., 2007. — *Grimmia dissimulata* nieuw voor België. *Dumortiera* **93**, in press.
- DE BEER D. & MARTENS C., 2007. — *Bryum dyffrynense* in Zeebrugge en Zwijndrecht, nieuw voor België. *Dumortiera* **91**: 25-26.
- DELLOGNE C.-H., 1883. — Flore cryptogamique de Belgique. *Annales de la Société Belge de Microscopie* **7**: 19-130.
- DE ROO R.T., HEDDERSON T.A. & SÖDERSTRÖM L., 2007. — Molecular insights into the phylogeny of the leafy liverwort family Lophoziaceae Cavers. *Taxon* **56**: 301-314.
- DE SLOOVER J.-L., 1970. — Que valent les collections de Pierre Troch? *Bull. Jard. Bot. Nat. Belg.* **40**: 19-21.
- DE SLOOVER J.-L. & STIEPERAERE H., 1999. — Le *Buxbaumia* récolté à Gentinnes par de Limminghe. *Dumortiera* **73**: 21-23.
- DE ZUTTERE P., 1993. — *Fissidens monguilloni* Thér. (Musci) espèce méconnue en Belgique et au Grand-Duché de Luxembourg. *Nowellia Bryologica* **3-4**: 1-4.
- DE ZUTTERE P. & POHL H., 1991. — Présence de deux hépatiques *Lophozia grandiretis* (Lindb. ex Kaal) Schiffn. et *Scapania praetervisa* Meyl. dans la partie ardennaise de l'Entre Sambre et Meuse. *Natura Mosana* **44**: 87-93.
- DE ZUTTERE P. & SOTIAUX A., 2007. — *Catascopium nigritum* (Hedw.) Brid. à la Baraque Michel (Prov. de Liège, Belgique). Que faut-il en penser ? *Nowellia Bryologica* **33**, in press.
- DE ZUTTERE P., SOTIAUX A. & POHL H., 2002. — Cartes provisoires de répartition de quelques mousses en Belgique. *Nowellia Bryologica* **23**: 24-40.
- ECCB (EUROPEAN COMMITTEE FOR THE CONSERVATION OF BRYOPHYTES), 1995. — *Red data book of European bryophytes*. Trondheim.

- ENGEL J.J. & BRAGGINS J.E., 2005. — Are *Mylia* and *Trabacellula* (Hepaticae) related? Unsuspected links revealed by cell wall morphology, with the transfer of *Mylia anomala* to a new genus (*Leiomylia* J.J. Engel and Braggins) of Jungermanniaceae. *Taxon* **54**: 665-680.
- ENGEL J.J. & SCHUSTER R.M., 1984. — An overview and evaluation of the genera Geocalycaceae subfamily Lophocoleoideae (Hepaticae). *Nova Hewigia* **39**: 385-463.
- FRAHM J.P., 2005. — New or interesting records of bryophytes from the Azores. *Tropical Bryology* **26**: 45-48.
- GREVEN H.C., 1995. — *Grimmia Hedw. (Grimmiaceae, Musci) in Europe*. Backhuys, Leiden.
- GREVEN H.C. & SOTIAUX A., 2003. — The boreal alpine bryophyte *Grimmia elongata* Kaulf. still present at Willerzie (Belgium) after 132 years. *Belg. J. Bot.* **136**: 165-166.
- GROLLE R. & LONG D.G., 2000. — An annotated check-list of the Hepaticae and Anthocerotae of Europe and Macaronesia. *J. Bryol.* **22**: 103-140.
- GROLLE R. & SO M.L., 2003. — *Riccia fruticulosa* O.F.Müll, 1782 and blue *Metzgeria* (Marchantiophyta) in Europe. *Bot. J. Linn. Soc.* **142**: 229-236.
- GUERRA J. & CANO M.J., 2000. — A taxonomic contribution on the European cleistocarpous species of Pottiaceae (Musci). *J. Bryol.* **22**: 91-97.
- HENTSCHEL J., ZÜNDORF H.-J., HELLWIG F.H., SCHÄFER-VERWIMP A. & HEINRICHS J., 2006. — Taxonomic studies in *Chiloscyphus* Corda (Jungermanniales: Lophocoleaceae) based on nrITS sequences and morphology. *Plant Syst. Evol.* **262**: 125-137.
- HEYLEN O. & WALRAEVENS A., 2001. — Spaanse Haarmuts in de Benelux: kabouter of 'zone skipper'? *Wielewaal* **67**: 50-54.
- HEYLEN O., JANSENS C. & GYSELS J., 2001. — Over de reanimatie van stekelhauwmossen (*Anthoceros caucasicus* Steph.) en andere hauwmossen. *Wielewaal* **67**: 13-14.
- HILL M.O., BELL N., BRUGGEMAN-NANNENGA M.A., BRUGUÉS M., CANO M.J., ENROTH J. et al., 2006. — An annotated checklist of the mosses of Europe and Macaronesia. *J. Bryol.* **28**: 198-267.
- KICKX J., 1867. — *Flore cryptogamique des Flandres. Tome premier*: 521p. Librairie H. Hoste, Gand, Librairie de J.B. Bailliére et fils. Paris.
- MUÑOZ J. & PANDO F., 2000. — *A world synopsis of the genus Grimmia (Musci, Grimmiaceae)*. Missouri Botanical Garden Press, St-Louis.
- RAEYMAEKERS G., 1981. — Bijdrage tot de kennis van de bladmossen van België I. *Philonotis, Pla-* giopus, *Breutelia, Catascopium*. *Dumortiera* **18**: 17-32.
- REINEKE G., SULT C., SCHMIDT C., LÜNTERBUSCH C. & DANIELS F.J.A., 2002. — *Polytrichum alpinum*, une mousse nouvelle pour la Belgique. *Dumortiera* **79**: 8-9.
- SCHUMACKER R. & SOTIAUX A., 2003. — *Rhytidadelphus subpinnatus*. In: BLOCKEEL T.L. (ed.), *New national and regional records*. *J. Bryol.* **25**: 217-221.
- SCHUMACKER R., DE ZUTTERE P. & LECLERCQ L., 1982. — *Anomodon rostratus* (Hedw.) Schimp. (Thuidiaceae), new for the Belgian bryoflora, in the south of the Ardenne massif (Rochehaut, prov. Luxembourg, Belgium). *J. Bryol.* **12**: 171-177.
- SCHUMACKER R., DE ZUTTERE P. & WERNER J., 1985. — Liste des bryophytes de Belgique et du Grand-Duché de Luxembourg. *Dumortiera* **31 suppl**: 28 pp.
- SHAW A.J. & ALLEN B., 2000. — Phylogenetic relationships, morphological incongruence, and geographic speciation in the Fontinalaceae (Bryophyta). *Mol. Phylogenet. Evol.* **16**: 225-237.
- SIEBEL H. & DURING H., 2006. — *Beknopte Mosflora van Nederland and België*. Stichting Uitgeverij KNNV, Utrecht.
- SIEBEL H.N., DURING H.J. & VAN MELICK H.M.H., 2005. — Veranderingen in de Standaardlijst van de Nederlandse blad-, lever- en hauwmossen (2005). *Buxbaumiella* **73**: 26-64.
- SMITH A.J.E., 2004. — *The Moss Flora of Britain and Ireland*. Cambridge University Press, Cambridge.
- SOTIAUX A. & SOTIAUX O., 1999. — *Ulota rehmannii* (Orthotrichaceae, Musci) nouveau pour la bryoflore belge dans le massif ardennais. *Belg. J. Bot.* **132**: 153-157.
- SOTIAUX A. & SOTIAUX O., 2002. — *Orthotrichum rogeri* (Orthotrichaceae, Musci) mousse épiphyte nouvelle pour la bryoflore belge, dans le district ardennais. Clé d'identification des *Orthotrichum* de Belgique. *Belg. J. Bot.* **135**: 27-31.
- SOTIAUX A. & SOTIAUX O., 2005. — *Tortula mucronifolia* (Pottiaceae, Musci), espèce boréo-montagnarde nouvelle pour la bryoflore belge dans la région de Soignies (province du Hainaut). *Dumortiera* **84**: 15-17.
- SOTIAUX A. & VANDERPOORTEN A., 2001. — Check-list of the bryophytes of Belgium. *Belg. J. Bot.* **134**: 97-120.
- SOTIAUX A. & VANDERPOORTEN A., 2004. — Catalogue, atlas commenté et mesures de conservation des bryophytes du bassin hydrographique de la Semois (Belgique, France). *Lejeunia* **175**: 1-47.

- SOTIAUX A., STIEPERAERE H. & SOTIAUX O., 1998a. — *Orthotrichum consimile* Mitt., an overlooked species in Europe? *J. Bryol.* **20**: 449-454.
- SOTIAUX A., SOTIAUX O. & PIERROT R.B., 1998b. — *Orthotrichum scanicum* (Orthotrichaceae, Musci) nouveau pour la bryoflore belge, dans le district ardennais. *Belg. J. Bot.* **131**: 23-27.
- SOTIAUX A., SOTIAUX O. & DUVIVIER J.P., 2002. — *Scapania calcicola*, hépatique boréo-montagnarde nouvelle pour la bryoflore belge, dans le district mosan. *Belg. J. Bot.* **135**: 88-91.
- SOTIAUX A., SOTIAUX O. & VANDERPOORTEN A., 2006. — Discovery of the rare liverwort *Haplomitrium hookeri* in Belgium: relictualism or long-distance dispersal? *Cryptogamie, Bryol.* **27**: 367-373.
- STIEPERAERE H., 2002. — *Anthoceros caucasicus* reeds in 1885 in de moerassen bij Geel. *Dumortiera* **79**: 25.
- STIEPERAERE H., 2007. — De mossen van enkele 'heide'-terreinen in het Provinciedomein Lippeveld-Bulskampveld (Beernem, West-Vlaanderen), n.a.v. de VWBL-excursie op 1 april 2006. *Muscillanea* **26**: 17-26.
- SZWEYKOWSKI J., BUCZKOWSKA K. & ODRZYKOSKI I.J., 2005. — *Conocephalum salebrosum* (Marchantiopsida, Conocephalaceae) – a new Holarctic liverwort species. *Plant Syst. Evol.* **253**: 133-158.
- VANDERPOORTEN A. & JACQUEMART A.L., 2004. — Evolutionary mode, tempo, and phylogenetic association of continuous morphological traits in the aquatic moss genus *Amblystegium*. *J. Evol. Biol.* **17**: 279-287.
- VANDERPOORTEN A., SOTIAUX A. & SOTIAUX O., 2001. — Integrating bryophytes into a forest management plan: lessons from grid-mapping in the forest of Soignes (Belgium). *Cryptogamie, Bryol.* **22**: 217-230.
- VANDERPOORTEN A., STIEPERAERE H. & KLEIN J.P., 1996. — Two rare European *Ephemerum* species: *E. cohaerens* rediscovered near Strasbourg and *E. stellatum* new for Belgium. *Belg. J. Bot.* **129**: 33-37.
- VANDERPOORTEN A., BOLES S. & SHAW A.J., 2003. — Patterns of molecular and morphological variation in *Leucobryum albidum*, *L. glaucum*, and *L. juniperoides* (Bryopsida). *Syst. Bot.* **28**: 651-656.
- VANDERPOORTEN A., COX C.J. & SHAW A.J., 2004. — Evolution of multiple paralogous adenosine kinase genes in the moss genus *Hygroamblystegium*: phylogenetic implications. *Mol. Phylogenet. Evol.* **31**: 505-516.
- VANDERPOORTEN A., SOTIAUX A. & ENGELS P., 2006. — A GIS-based model of the distribution of the rare liverwort *Aneura maxima* at the landscape scale for an improved assessment of its conservation status. *Biodiv. Conserv.* **15**: 829-838.
- WERNER J., 2003. — Liste rouge des bryophytes du Luxembourg. *Ferrantia* **35**: 1-71.
- WERNER O., ROS R.M. & GRUNDMANN M., 2005a. — Molecular phylogeny of Trichostomoideae (Pottiaceae, Bryophyta) based on nrITS sequence data. *Taxon* **54**: 361-368.
- WERNER O., JIMENEZ J.A., ROS R.M., CANO M.J. & GUERRA J., 2005b. — Preliminary investigation of the systematics of *Didymodon* (Pottiaceae, Musci) based on nrITS sequence data. *Syst. Bot.* **30**: 461-470.
- ZANDER R.H., 1993. — *Genera of the Pottiaceae: mosses of harsh environments*. Buffalo Society of Natural Sciences, New York.

APPENDIX

Checklist of the bryophytes of Belgium. O: occurrence within each region (F = Flanders, W = Wallonia, B = Brussels-Capital Region). Species names in bold denote species likely to be extinct because no occurrence has been reported in Belgium since 1980 or because the only known locality has been destroyed.

Taxon	O	O	O
Hornworts			
<i>Anthoceros agrestis</i> Paton	F	W	
<i>Anthoceros caucasicus</i> Steph.	F		
<i>Anthoceros punctatus</i> L.	F	W	
<i>Phaeoceros carolinianus</i> (Michx.) Prosk.	F	W	
<i>Phaeoceros laevis</i> (L.) Prosk.	F		

Liverworts

<i>Anastrophyllum minutum</i> (Schreb.) R.M.Schust.	F	W
<i>Aneura maxima</i> (Schiffn.) Steph.	F	W
<i>Aneura pinguis</i> (L.) Dumort.	F	W
<i>Apometzgeria pubescens</i> (Schrank) Kuwah.		W
<i>Barbilophozia attenuata</i> (Mart.) Loeske	F	W
<i>Barbilophozia barbata</i> (Schmidel ex Schreb.) Loeske	F	W
<i>Barbilophozia floerkei</i> (F.Weber & D.Mohr) Loeske	F	W
<i>Barbilophozia hatcheri</i> (A.Evans) Loeske	F	W
<i>Barbilophozia kunzeana</i> (Huebener) Müll.Frib.	F	W
<i>Bazzania flaccida</i> (Dumort.) Grolle		W
<i>Bazzania trilobata</i> (L.) S.F.Gray	F	W
<i>Blasia pusilla</i> L.	F	W
<i>Blepharostoma trichophyllum</i> (L.) Dumort.	F	W
<i>Calypogeia arguta</i> Nees & Mont.	F	W
<i>Calypogeia azurea</i> Stotler & Crotz	F	W
<i>Calypogeia fissa</i> (L.) Raddi	F	W
<i>Calypogeia integrifistula</i> Steph.	F	W
<i>Calypogeia muelleriana</i> (Schiffn.) Müll.Frib.	F	W
<i>Calypogeia neesiana</i> (C.Mass. & Carestia.) Müll.Frib.	F	W
<i>Calypogeia sphagnicola</i> (Arnell. & J.Perss.) Warnst. & Loeske	F	
<i>Calypogeia suecica</i> (Arnell & J.Perss.) Müll.Frib.		W
<i>Cephalozia bicuspidata</i> (L.) Dumort.	F	W
<i>Cephalozia catenulata</i> (Huebener) Lindb.		W
<i>Cephalozia connivens</i> (Dicks.) Lindb.	F	W
<i>Cephalozia lunulifolia</i> (Dumort.) Dumort.	F	W
<i>Cephalozia macrostachya</i> Kaal.	F	
<i>Cephalozia pleniceps</i> (Austin) Lindb.	F	W
<i>Cephaloziella baumgartneri</i> Schiffn.		W
<i>Cephaloziella divaricata</i> (Sm.) Schiffn.	F	W
<i>Cephaloziella elachista</i> (J.B.Jack ex Gottsche & Rabenh.) Schiffn.	F	W
<i>Cephaloziella hampeana</i> (Nees) Schiffn.	F	W
<i>Cephaloziella rubella</i> (Nees) Warnst.	F	W
<i>Cephaloziella stellulifera</i> (Taylor ex Spruce) Schiffn.	F	W
<i>Chiloscyphus pallescens</i> (Ehrh. ex Hoffm.) Dumort.	F	W
<i>Chiloscyphus polyanthos</i> (L.) Corda	F	W
<i>Cladopodiella fluitans</i> (Nees) H.Buch.	F	W
<i>Cladopodiella francisci</i> (Hook.) Jörg.	F	W
<i>Cololejeunea calcarea</i> (Libert) Schiffn.		W
<i>Cololejeunea minutissima</i> (Sm.) Schiffn.	F	W
<i>Cololejeunea rossettiana</i> (C.Massal.) Schiffn.		W
<i>Conocephalum conicum</i> (L.) Dumort.	F	W
<i>Conocephalum salebrosum</i> Szweykowski, Buczkowska & Odrzykoski	F	W
<i>Diplophyllum albicans</i> (L.) Dumort.	F	W
<i>Diplophyllum obtusifolium</i> (Hook.) Dumort.	F	W
<i>Douinia ovata</i> (Dicks.) H.Buch		W
<i>Fossombronia foveolata</i> Lindb.	F	W
<i>Fossombronia incurva</i> Lindb.	F	W
<i>Fossombronia pusilla</i> (L.) Nees	F	W
<i>Fossombronia wondraczekii</i> (Corda) Lindb.	F	W
<i>Frullania dilatata</i> (L.) Dumort.	F	W
<i>Frullania fragilifolia</i> (Taylor) Gottsche et al.		W
<i>Frullania tamarisci</i> (L.) Dumort.	F	W
<i>Gymnocolea inflata</i> (Huds.) Dumort.	F	W
<i>Haplomitrium hookeri</i> (Sm.) Nees		W
<i>Harpanthus flotovianus</i> (Nees) Nees	F	
<i>Harpanthus scutatus</i> (F.Weber & D.Mohr) Spruce		W
<i>Jamesoniella autumnalis</i> (DC.) Steph.		W

<i>Jungermannia atrovirens</i> Dumort.		W
<i>Jungermannia caespiticia</i> Lindenb.	F	W
<i>Jungermannia exsertifolia</i> Steph.		W
<i>Jungermannia gracillima</i> Sm.	F	W
<i>Jungermannia hyalina</i> Lyell	F	W
<i>Jungermannia leiantha</i> Grolle		W
<i>Jungermannia obovata</i> Nees		W
<i>Jungermannia paroica</i> (Schiffn.) Grolle		W
<i>Jungermannia pumila</i> With.		W
<i>Jungermannia sphaerocarpa</i> Hook.		W
<i>Jungermannia subelliptica</i> (Lindb. ex Kaal.) Levier		W
<i>Kurzia pauciflora</i> (Dicks.) Grolle	F	W
<i>Kurzia sylvatica</i> (A.Evans) Grolle	F	W
<i>Kurzia trichoclados</i> (Müll.Frib.) Grolle		W
<i>Leiocolea badensis</i> (Gottsc.) Jörg.	F	W
<i>Leiocolea collaris</i> (Nees) Schljakov	F	W
<i>Leiocolea heterocolpos</i> (Thed. ex Hartm.) H.Buch		W
<i>Leiocolea turbinata</i> (Raddi) H.Buch		W
<i>Lejeunea cavifolia</i> (Ehrh.) Lindb.	F	W
<i>Lejeunea lamacerina</i> (Steph.) Schiffn.		W
<i>Lepidozia reptans</i> (L.) Dumort.	F	W
<i>Lophocolea bidentata</i> (L.) Dumort.	F	W
<i>Lophocolea heterophylla</i> (Schrad.) Dumort.	F	W
<i>Lophocolea minor</i> Nees	F	W
<i>Lophocolea semiteres</i> (Lehm.) Mitt.	F	W
<i>Lophozia bicrenata</i> (Schmidel ex Hoffm.) Dumort.	F	W
<i>Lophozia capitata</i> (Hook.) Macoun	F	W
<i>Lophozia excisa</i> (Dicks.) Dumort.	F	W
<i>Lophozia grandiretis</i> (Lindb. ex Kaal.) Schiffn.	F	W
<i>Lophozia incisa</i> (Schrad.) Dumort.	F	W
<i>Lophozia longidens</i> (Lindb.) Macoun		W
<i>Lophozia longiflora</i> (Nees) Schiffn.		W
<i>Lophozia obtusa</i> (Lindb.) A.Evans		W
<i>Lophozia perssonii</i> H.Buch & S.W.Arnell	F	W
<i>Lophozia sudetica</i> (Nees ex Huebener) Grolle		W
<i>Lophozia ventricosa</i> (Dicks.) Dumort.	F	W
<i>Lophozia wenzelii</i> (Nees) Steph.		W
<i>Lunularia cruciata</i> (L.) Lindb.	F	W
<i>Marchantia polymorpha</i> L.	F	W
<i>Marsupella emarginata</i> (Ehrh.) Dumort.	F	W
<i>Marsupella funkii</i> (F.Weber & D.Mohr) Dumort.	F	W
<i>Marsupella sphacelata</i> (Gies. ex Lindenb.) Dumort.		W
<i>Marsupella sprucei</i> (Limpr.) Bernet		W
<i>Metzgeria conjugata</i> Lindb.		W
<i>Metzgeria furcata</i> (L.) Dumort.	F	W
<i>Metzgeria temperata</i> Kuwah.	F	W
<i>Metzgeria violacea</i> (Ach.) Dumort.	F	W
<i>Microlejeunea ulicina</i> (Taylor) A.Evans	F	W
<i>Moerckia hibernica</i> (Hook.) Gottsc.	F	
<i>Mylia anomala</i> (Hook.) Gray	F	W
<i>Mylia taylorii</i> (Hook.) Gray		W
<i>Nardia compressa</i> (Hook.) Gray	F	W
<i>Nardia geoscyphus</i> (De Not.) Lindb.	F	W
<i>Nardia insecta</i> Lindb.		W
<i>Nardia scalaris</i> Gray	F	W
<i>Nowellia curvifolia</i> (Dicks.) Mitt.	F	W
<i>Odontoschisma denudatum</i> (Mart.) Dumort.	F	W
<i>Odontoschisma sphagni</i> (Dicks.) Dumort.	F	W

<i>Pallavicinia lyellii</i> (Hook.) Carruth.	F	W
<i>Pedinophyllum interruptum</i> (Nees) Kaal.		W
<i>Pellia endiviifolia</i> (Dicks.) Dumort.	F	W
<i>Pellia epiphylla</i> (L.) Corda	F	W
<i>Pellia neesiana</i> (Gottsche) Limpr.	F	W
<i>Plagiochila asplenoides</i> (L. emend. Taylor) Dumort.	F	W
<i>Plagiochila bifaria</i> (Sw.) Lindenb.		W
<i>Plagiochila poreloides</i> (Torrey ex Nees) Lindenb.	F	W
<i>Plagiochila spinulosa</i> (Dicks.) Dumort.		W
<i>Porella arboris-vitae</i> (With.) Grolle		W
<i>Porella cordaeana</i> (Huebener) Moore		W
<i>Porella platyphylla</i> (L.) Pfeiff.	F	W
<i>Preissia quadrata</i> (Scop.) Nees	F	W
<i>Ptilidium ciliare</i> (L.) Hampe	F	W
<i>Ptilidium pulcherrimum</i> (Weber) Vainio	F	W
<i>Radula complanata</i> (L.) Dumort.	F	W
<i>Radula lindenbergiana</i> Gottsche ex C.Hartm.		W
<i>Reboulia hemisphaerica</i> (L.) Raddi	F	W
<i>Riccardia chamedryfolia</i> (With.) Grolle	F	W
<i>Riccardia incurvata</i> Lindb.	F	W
<i>Riccardia latifrons</i> (Lindb.) Lindb.	F	W
<i>Riccardia multifida</i> (L.) Gray	F	W
<i>Riccia beyrichiana</i> Hampe ex Lehm.	F	W
<i>Riccia bifurca</i> Hoffm.	F	W
<i>Riccia canaliculata</i> Hoffm.	F	W
<i>Riccia cavernosa</i> Hoffm.	F	W
<i>Riccia ciliata</i> Hoffm.	F	W
<i>Riccia ciliifera</i> Link ex Lindenb.		W
<i>Riccia fluitans</i> L.	F	W
<i>Riccia glauca</i> L.	F	W
<i>Riccia huebeneriana</i> Lindenb.	F	W
<i>Riccia rhenana</i> Lorb. ex Müll.Frib.		W
<i>Riccia sorocarpa</i> Bisch.	F	W
<i>Riccia subbifurca</i> Warnst. ex Croz.	F	W
<i>Riccia warnstorffii</i> Limpr. ex Warnst.	F	W
<i>Ricciocarpus natans</i> (L.) Corda	F	W
<i>Scapania aequiloba</i> (Schwägr.) Dumort.		W
<i>Scapania aspera</i> Bernet & M.Bern.		W
<i>Scapania calcicola</i> (Arnell & J.Perss.) Ingham		W
<i>Scapania compacta</i> (A.Roth) Dumort.	F	W
<i>Scapania curta</i> (Mart.) Dumort.	F	W
<i>Scapania cuspiduligera</i> (Nees) K.Müll.		W
<i>Scapania irrigua</i> (Nees) Nees	F	W
<i>Scapania lingulata</i> H.Buch		W
<i>Scapania mucronata</i> H.Buch		W
<i>Scapania nemorea</i> (L.) Grolle	F	W
<i>Scapania paludicola</i> Loeske & Mül.Frib.	F	W
<i>Scapania scandica</i> (Arnell & H.Buch) Macvicar	F	W
<i>Scapania undulata</i> (L.) Dumort.	F	W
<i>Southbya nigrella</i> (De Not.) Henriq.		W
<i>Sphaerocarpos michelii</i> Bellardi	F	W
<i>Sphaerocarpos texanus</i> Austin	F	W
<i>Targionia hypophylla</i> L.	F	W
<i>Tetralophozia setiformis</i> (Ehrh.) Schljakov		W
<i>Trichocolea tomentella</i> (Ehrh.) Dumort.	F	W
<i>Tritomaria exsecta</i> (Schmidel) Loeske		W
<i>Tritomaria exsectiformis</i> (Breidl.) Loeske	F	W
<i>Tritomaria quinquedentata</i> (Huds.) H.Buch	F	W

Mosses

<i>Abietinella abietina</i> (Hedw.) M.Fleisch.	F	W	
<i>Acaulon muticum</i> (Hedw.) Müll.Hal.	F	W	B
<i>Acaulon triquetrum</i> (Spruce) Müll.Hal.		W	
<i>Aloina aloides</i> (Koch ex Schultz) Kindb.	F	W	B
<i>Aloina ambigua</i> (Bruch & Schimp.) Limpr.	F	W	B
<i>Aloina rigida</i> (Hedw.) Limpr.	F	W	
<i>Amblystegium confervoides</i> (Brid.) Schimp.		W	
<i>Amblystegium radicale</i> (P.Beauv.) Schimp.	F	W	
<i>Amblystegium serpens</i> (Hedw.) Schimp.	F	W	B
<i>Amphidium mougeotii</i> (Schimp.) Schimp.		W	
<i>Andreaea rothii</i> F.Weber & D.Mohr		W	
<i>Andreaea rupestris</i> Hedw.		W	
<i>Anomobryum concinnatum</i> (Spruce) Lindb.		W	
<i>Anomobryum julaceum</i> (Schrad. ex P.Gaertn. et al.) Schimp.		W	
<i>Anomodon attenuatus</i> (Hedw.) Huebener	F	W	
<i>Anomodon longifolius</i> (Schleich. ex Brid.) Hartm.		W	
<i>Anomodon rostratus</i> (Hedw.) Schimp.		W	
<i>Anomodon viticulosus</i> (Hedw.) Hook. & Taylor	F	W	B
<i>Antitrichia curtipendula</i> (Hedw.) Brid.	F	W	
<i>Archidium alternifolium</i> (Hedw.) Mitt.	F	W	
<i>Atrichum angustatum</i> (Brid.) Bruch & Schimp.		W	
<i>Atrichum tenellum</i> (Röhl.) Bruch & Schimp.	F	W	
<i>Atrichum undulatum</i> (Hedw.) P.Beauv.	F	W	B
<i>Aulacomnium androgynum</i> (Hedw.) Schwägr.	F	W	B
<i>Aulacomnium palustre</i> (Hedw.) Schwägr.	F	W	
<i>Barbula convoluta</i> Hedw.	F	W	B
<i>Barbula unguiculata</i> Hedw.	F	W	B
<i>Bartramia halleriana</i> Hedw.		W	
<i>Bartramia ithyphylla</i> Brid.		W	
<i>Bartramia pomiformis</i> Hedw.	F	W	
<i>Brachytheciastrum velutinum</i> (Hedw.) Ignatov & Huttunen	F	W	B
<i>Brachythecium albicans</i> (Hedw.) Schimp.	F	W	B
<i>Brachythecium campestre</i> (Müll.Hal.) Schimp.		W	
<i>Brachythecium glareosum</i> (Bruch ex Spruce) Schimp.	F	W	B
<i>Brachythecium laetum</i> (Brid.) Schimp.		W	
<i>Brachythecium mildeanum</i> (Schimp.) Schimp.	F	W	B
<i>Brachythecium rivulare</i> Schimp.	F	W	B
<i>Brachythecium rutabulum</i> (Hedw.) Schimp.	F	W	B
<i>Brachythecium salebrosum</i> (Hoffm. ex F.Weber & D.Mohr) Schimp.	F	W	B
<i>Brachythecium tommasinii</i> (Sendtn. ex Boulay) Ignatov & Huttunen		W	
<i>Breidleria pratinensis</i> (W.D.J.Koch ex Spruce) Loeske		W	
<i>Breutelia chrysocoma</i> (Hedw.) Lindb.		W	
<i>Bryoerythrophyllum ferruginascens</i> (Stirt.) Giacom.		W	
<i>Bryoerythrophyllum recurvirostrum</i> (Hedw.) P.C.Chen	F	W	B
<i>Bryum algovicum</i> Sendtn. ex Müll.Hal.	F	W	
<i>Bryum alpinum</i> Huds. ex With.	F	W	
<i>Bryum archangelicum</i> Bruch & Schimp.	F	W	
<i>Bryum argenteum</i> Hedw.	F	W	B
<i>Bryum bornholmense</i> Wink. & R.Ruthe	F	W	
<i>Bryum caespiticium</i> Hedw.	F	W	B
<i>Bryum canariense</i> Brid.		W	
<i>Bryum capillare</i> Hedw.	F	W	B
<i>Bryum creberrimum</i> Taylor	F	W	
<i>Bryum cyclophyllum</i> (Schwägr.) Bruch & Schimp.	F		
<i>Bryum demareitianum</i> Arts	F		
<i>Bryum dichotomum</i> Hedw.	F	W	B
<i>Bryum donianum</i> Grev.		W	

<i>Bryum dyffrynense</i> Holyoak	F	
<i>Bryum elegans</i> Nees		W
<i>Bryum funckii</i> Schwägr.		W
<i>Bryum gemmiferum</i> R.Wilczek & Demaret	F	W
<i>Bryum gemmilucens</i> R.Wilczek & Demaret	F	W
<i>Bryum gemmiparum</i> De Not.		W
<i>Bryum intermedium</i> (Brid.) Blandow	F	W
<i>Bryum klinggraeffii</i> Schimp.	F	W
<i>Bryum knowltonii</i> Barnes	F	
<i>Bryum kunzei</i> Hornsch.		W
<i>Bryum mildeanum</i> Jur.	F	W
<i>Bryum moravicum</i> Podp.	F	W
<i>Bryum muehlenbeckii</i> Bruch & Schimp.		B
<i>Bryum neodamense</i> Itzigs.	F	
<i>Bryum pallens</i> Sw. ex anon.	F	W
<i>Bryum pallescens</i> Schleich. ex Schwägr.	F	W
<i>Bryum pseudotriquetrum</i> (Hedw.) P.Gaertn. et al.	F	W
<i>Bryum radiculosum</i> Brid.	F	W
<i>Bryum rubens</i> Mitt.	F	W
<i>Bryum ruderale</i> Crundw. & Nyholm	F	W
<i>Bryum sauteri</i> Bruch & Schimp.	F	W
<i>Bryum subapiculatum</i> Hampe	F	W
<i>Bryum tenuisetum</i> Limpr.	F	B
<i>Bryum torquescens</i> Bruch & Schimp.		W
<i>Bryum turbinatum</i> (Hedw.) Turner		W
<i>Bryum uliginosum</i> (Brid.) Bruch & Schimp.	F	W
<i>Bryum violaceum</i> Crundw. & Nyholm	F	W
<i>Bryum warneum</i> (Röhl.) Brid.	F	
<i>Bryum weigelii</i> Spreng.	F	W
<i>Buxbaumia aphylla</i> Hedw.	F	W
<i>Calliergon cordifolium</i> (Hedw.) Kindb.	F	W
<i>Calliergon giganteum</i> (Schimp.) Kindb.	F	W
<i>Calliergonella cuspidata</i> (Hedw.) Loeske	F	W
<i>Calliergonella lindbergii</i> (Mitt.) Hedenäs	F	B
<i>Campyliadelphus chrysophyllus</i> (Brid.) R.S.Chopra	F	W
<i>Campyliadelphus elodes</i> (Lindb.) Kanda	F	W
<i>Campylium protensum</i> (Brid.) Kindb.	F	W
<i>Campylium stellatum</i> (Hedw.) Lange & C.E.O.Jensen	F	W
<i>Campylophyllum calcareum</i> (Crundw. & Nyholm) Hedenäs	F	W
<i>Campylopus brevipilus</i> Bruch & Schimp.	F	
<i>Campylopus flexuosus</i> (Hedw.) Brid.	F	W
<i>Campylopus fragilis</i> (Brid.) Bruch & Schimp.	F	W
<i>Campylopus introflexus</i> (Hedw.) Brid.	F	B
<i>Campylopus pilifer</i> Brid.	F	
<i>Campylopus pyriformis</i> (Schultz) Brid.	F	W
<i>Campylopus subulatus</i> Schimp. ex Milde		W
<i>Campylostelium saxicola</i> (F.Weber & D.Mohr) Bruch & Schimp.		W
<i>Catoscopium nigritum</i> (Hedw.) Brid.	F	W
<i>Ceratodon conicus</i> (Hampe) Lindb.		W
<i>Ceratodon purpureus</i> (Hedw.) Brid.	F	W
<i>Cinclidium stygium</i> Sw.		B
<i>Cinclidotus aquaticus</i> (Hedw.) Bruch & Schimp.		W
<i>Cinclidotus danubicus</i> Schiffn. & Baumgartner	F	W
<i>Cinclidotus fontinaloides</i> (Hedw.) P.Beauv.	F	W
<i>Cinclidotus riparius</i> (Host ex Brid.) Arn.	F	B
<i>Cirriphyllum crassinervium</i> (Taylor) Loeske & M.Fleisch.	F	W
<i>Cirriphyllum piliferum</i> (Hedw.) Grout	F	B

<i>Cleistocarpidium palustre</i> (Bruch & Schimp)	Ochyra & Bednarek-Ochyra	F	W	
<i>Climaciumpendroides</i> (Hedw.) F.Weber & D.Mohr		F	W	B
<i>Conardia compacta</i> (Müll.Hal.) H.Rob.			W	
<i>Coscinodon cribrosus</i> (Hedw.) Spruce			W	
<i>Cratoneuron filicinum</i> (Hedw.) Spruce		F	W	B
<i>Crossidium squamiferum</i> (Viv.) Jur.			W	
<i>Cryphaea heteromalla</i> (Hedw.) D.Mohr		F	W	B
<i>Ctenidium molluscum</i> (Hedw.) Mitt.		F	W	B
<i>Cynodontium bruntonii</i> (Sm.) Bruch & Schimp.			W	
<i>Cynodontium polycarpon</i> (Hedw.) Schimp.			W	
<i>Cynodontium strumiferum</i> (Hedw.) Lindb.			W	
<i>Dalytrichia fragilifolia</i> (Bizot & J.Roux) F.Lara			W	
<i>Dalytrichia mucronata</i> (Brid.) Broth.			W	
<i>Dichodontium flavescens</i> (Dicks.) Lindb.			W	
<i>Dichodontium palustre</i> (Dicks.) M.Stech			W	
<i>Dichodontium pellucidum</i> (Hedw.) Schimp.			W	B
<i>Dicranella cerviculata</i> (Hedw.) Schimp.		F	W	B
<i>Dicranella heteromalla</i> (Hedw.) Schimp.		F	W	B
<i>Dicranella rufescens</i> (Dicks.) Schimp.		F	W	B
<i>Dicranella schreberiana</i> (Hedw.) Dixon		F	W	B
<i>Dicranella staphylina</i> H.Whitehouse		F	W	B
<i>Dicranella subulata</i> (Hedw.) Schimp.			W	
<i>Dicranella varia</i> (Hedw.) Schimp.		F	W	B
<i>Dicranodontium denudatum</i> (Brid.) E.Britton			W	
<i>Dicranoweisia cirrata</i> (Hedw.) Lindb.		F	W	B
<i>Dicranum bonjeanii</i> De Not.		F	W	
<i>Dicranum flagellare</i> Hedw.		F	W	
<i>Dicranum fulvum</i> Hook.			W	
<i>Dicranum fuscescens</i> Sm.			W	
<i>Dicranum majus</i> Sm.		F	W	B
<i>Dicranum montanum</i> Hedw.		F	W	B
<i>Dicranum polysetum</i> Sw. ex anon.		F	W	
<i>Dicranum scoparium</i> Hedw.		F	W	B
<i>Dicranum spuriu</i> Hedw.		F	W	
<i>Dicranum tauricum</i> Sapjegin		F	W	B
<i>Dicranum undulatum</i> Schrad. ex Brid.			W	
<i>Dicranum viride</i> (Sull. & Lesq.) Lindb.			W	
<i>Didymodon acutus</i> (Brid.) K.Saito		F	W	
<i>Didymodon australasiae</i> (Hook. & Grev.) R.H.Zander		F	W	
<i>Didymodon cordatus</i> Jur.			W	
<i>Didymodon fallax</i> (Hedw.) R.H.Zander		F	W	B
<i>Didymodon ferrugineus</i> (Schimp. ex Besch.) M.O.Hill		F	W	
<i>Didymodon insulanus</i> (De Not.) M.O.Hill		F	W	B
<i>Didymodon luridus</i> Hornsch.		F	W	B
<i>Didymodon nicholsonii</i> Culm.			W	
<i>Didymodon rigidulus</i> Hedw.		F	W	B
<i>Didymodon sinuosus</i> (Mitt.) Delogne		F	W	B
<i>Didymodon spadiceus</i> (Mitt.) Limpr.		F	W	B
<i>Didymodon tophaceus</i> (Brid.) Lisa		F	W	B
<i>Didymodon vinealis</i> (Brid.) R.H.Zander		F	W	B
<i>Diphyscium foliosum</i> (Hedw.) D.Mohr		F	W	
<i>Discelium nudum</i> (Dicks.) Brid.			W	
<i>Distichium capillaceum</i> (Hedw.) Bruch & Schimp.		F	W	
<i>Distichium inclinatum</i> (Hedw.) Bruch & Schimp.			W	
<i>Ditrichum flexicaule</i> (Schwägr.) Hampe		F	W	
<i>Ditrichum gracile</i> (Mitt.) Kuntze		F	W	
<i>Ditrichum heteromallum</i> (Hedw.) E.Britton		F	W	B

<i>Ditrichum lineare</i> (Sw.) Lindb.	F	W
<i>Ditrichum pallidum</i> (Hedw.) Hampe	F	W
<i>Ditrichum pusillum</i> (Hedw.) Hampe	F	W
<i>Drepanocladus aduncus</i> (Hedw.) Warnst.	F	W
<i>Drepanocladus polygamus</i> (Schimp.) Hedenäs	F	W
<i>Drepanocladus sendtneri</i> (Schimp. ex H.Müll.) Warnst.	F	W
<i>Encalypta ciliata</i> Hedw.		W
<i>Encalypta streptocarpa</i> Hedw.	F	W
<i>Encalypta vulgaris</i> Hedw.	F	W
<i>Entodon concinnus</i> (De Not.) Paris	F	W
<i>Entosthodon fascicularis</i> (Hedw.) Müll.Hal	F	W
<i>Entosthodon muhlenbergii</i> (Turner) Fife		W
<i>Entosthodon obtusus</i> (Hedw.) Lindb.	F	W
<i>Entosthodon pulchellus</i> (H.Philib.) Brugués		W
<i>Ephemerum cohaerens</i> (Hedw.) Hampe		W
<i>Ephemerum minutissimum</i> Lindb.	F	W
<i>Ephemerum recurvifolium</i> (Dicks.) Boulay		W
<i>Ephemerum serratum</i> (Hedw.) Hampe	F	W
<i>Ephemerum sessile</i> (Bruch) Müll.Hal.		W
<i>Ephemerum stellatum</i> H.Philib.		B
<i>Eucladium verticillatum</i> (With.) Bruch & Schimp.	F	W
<i>Eurhynchiastrum pulchellum</i> (Hedw.) Ignatov & Huttunen	F	W
<i>Eurhynchium angustirete</i> (Broth.) T.J.Kop.		W
<i>Eurhynchium striatum</i> (Hedw.) Schimp.	F	W
<i>Fissidens adianthoides</i> Hedw.	F	W
<i>Fissidens arnoldii</i> R.Ruthe		W
<i>Fissidens bryoides</i> Hedw.	F	W
<i>Fissidens celticus</i> Paton		W
<i>Fissidens crassipes</i> Wilson ex Bruch & Schimp.	F	W
<i>Fissidens dubius</i> P.Beauv.	F	W
<i>Fissidens exilis</i> Hedw.	F	W
<i>Fissidens fontanii</i> (Bach.Pyl.) Steud.	F	W
<i>Fissidens gracilifolius</i> Brugg.-Nann. & Nyholm	F	W
<i>Fissidens monguilloni</i> Thér.	F	W
<i>Fissidens osmundoides</i> Hedw.	F	W
<i>Fissidens pusillus</i> (Wilson) Milde	F	W
<i>Fissidens rivularis</i> (Spruce) Schimp.		W
<i>Fissidens rufulus</i> Bruch & Schimp.	F	W
<i>Fissidens taxifolius</i> Hedw.	F	W
<i>Fissidens viridulus</i> (Sw. ex anon.) Wahlenb.	F	W
<i>Fontinalis antipyretica</i> Hedw.	F	W
<i>Fontinalis squamosa</i> Hedw.		W
<i>Funaria hygrometrica</i> Hedw.	F	W
<i>Grimmia atrata</i> Miel. ex Hornsch.		W
<i>Grimmia caespiticia</i> (Brid.) Jur.		W
<i>Grimmia crinita</i> Brid.	F	W
<i>Grimmia decipiens</i> (Schultz) Lindb.		W
<i>Grimmia dissimulata</i> E.Maier	F	?
<i>Grimmia elongata</i> Kaulf.		W
<i>Grimmia hartmanii</i> Schimp.		W
<i>Grimmia laevigata</i> (Brid.) Brid.		W
<i>Grimmia lisae</i> De Not.		W
<i>Grimmia longirostris</i> Hook.		W
<i>Grimmia montana</i> Bruch & Schimp.		W
<i>Grimmia muehlenbeckii</i> Schimp.		W
<i>Grimmia orbicularis</i> Bruch ex Wilson		W
<i>Grimmia ovalis</i> (Hedw.) Lindb.		W

	F	W	B
<i>Grimmia pulvinata</i> (Hedw.) Sm.			
<i>Grimmia ramondii</i> (Lam. & DC.) Margad.		W	
<i>Grimmia tergestina</i> Tomm. ex Bruch & Schimp.		W	
<i>Grimmia torquata</i> Drumm.		W	
<i>Grimmia trichophylla</i> Grev.	F	W	
<i>Gymnostomum aeruginosum</i> Sm.		W	
<i>Gymnostomum calcareum</i> Nees & Hornsch.		W	
<i>Gymnostomum viridulum</i> Brid.		W	
<i>Gyroweisia tenuis</i> (Hedw.) Schimp.	F	W	B
<i>Habrodon perpusillus</i> (De Not.) Lindb.		W	
<i>Hageniella micans</i> (Mitt.) B.C.Tan & Y.Jia		W	
<i>Hamatocaulis vernicosus</i> (Mitt.) Hedenäs	F	W	
<i>Hedwigia ciliata</i> (Hedw.) P.Beauv.	F	W	
<i>Hedwigia stellata</i> Hedenäs		W	
<i>Helodium blandowii</i> (F.Weber & D.Mohr.) Warnst.	F		
<i>Hennediella heimii</i> (Hedw.) R.H.Zander	F	W	
<i>Herzogiella seligeri</i> (Brid.) Z.Iwats.	F	W	B
<i>Heterocladium flaccidum</i> (Schimp.) A.J.E.Smith		W	
<i>Heterocladium heteropterum</i> (Brid.) Schimp.	F	W	
<i>Homalia trichomanoides</i> (Hedw.) Brid.	F	W	B
<i>Homalothecium lutescens</i> (Hedw.) H.Rob.	F	W	B
<i>Homalothecium sericeum</i> (Hedw.) Schimp.	F	W	B
<i>Homomallium incurvatum</i> (Schrad. ex Brid.) Loeske		W	
<i>Hookeria lucens</i> (Hedw.) Sm.	F	W	
<i>Hygroamblystegium fluviatile</i> (Hedw.) Loeske	F	W	
<i>Hygroamblystegium humile</i> (P.Beauv.) Vanderp., Goffinet & Hedenäs	F	W	B
<i>Hygroamblystegium tenax</i> (Hedw.) Jenn.	F	W	B
<i>Hygroamblystegium varium</i> (Hedw.) Mönk.	F	W	B
<i>Hygrohypnum eugyrium</i> (Schimp.) Broth.		W	
<i>Hygrohypnum luridum</i> (Hedw.) Jenn.	F	W	B
<i>Hygrohypnum ochraceum</i> (Turner ex Wilson) Loeske	F	W	
<i>Hylocomiastrum pyrenaeicum</i> (Spruce) M.Fleisch.		W	
<i>Hylocomiastrum umbratum</i> (Hedw.) M.Fleisch.		W	
<i>Hylocomium splendens</i> (Hedw.) Schimp.	F	W	
<i>Hymenostylium recurvirostrum</i> (Hedw.) Dixon		W	
<i>Hyocomium armoricum</i> (Brid.) Wijk & Margad.		W	
<i>Hypnum andoi</i> A.J.E.Smith		W	
<i>Hypnum cupressiforme</i> Hedw.	F	W	B
<i>Hypnum imponens</i> Hedw.	F	W	
<i>Hypnum jutlandicum</i> Holmen & E.Warncke	F	W	B
<i>Hypnum pallescens</i> (Hedw.) P.Beauv.	F	W	
<i>Isothecium alopecuroides</i> (Lam. ex Dubois) Isov.	F	W	B
<i>Isothecium holtii</i> Kindb.		W	
<i>Isothecium myosuroides</i> Brid.	F	W	B
<i>Kindbergia praelonga</i> (Hedw.) Ochyra	F	W	B
<i>Leptobarbula berica</i> (De Not.) Schimp.	F	W	
<i>Leptobryum pyriforme</i> (Hedw.) Wilson	F	W	
<i>Leptodictyum riparium</i> (Hedw.) Warnst.	F	W	B
<i>Leptodontium flexifolium</i> (Dicks.) Hampe	F	W	
<i>Leptodontium gemmascens</i> (Mitt.) Braithw.	F	W	
<i>Leskeia polycarpa</i> Hedw.	F	W	B
<i>Leucobryum glaucum</i> (Hedw.) Ångstr.	F	W	B
<i>Leucodon sciuroides</i> (Hedw.) Schwägr.	F	W	
<i>Loeskeobryum brevirostre</i> (Brid.) M.Fleisch.	F	W	
<i>Meesia triquetra</i> (L. ex Jolycl.) Ångstr.		W	
<i>Microbryum curvicollum</i> (Hedw.) R.H.Zander	F	W	
<i>Microbryum davallianum</i> (Sm.) R.H.Zander	F	W	B

<i>Microbryum floerkeanum</i> (F.Weber & D.Mohr) Schimp.	F	W
<i>Microbryum rectum</i> (With.) R.H.Zander		W
<i>Microbryum starckeanaum</i> (Hedw.) R.H.Zander	F	W
<i>Micromitrium tenerum</i> (Bruch & Schimp.) Crosby	F	
<i>Mielichhoferia mielichhoferiana</i> (Funck) Loeske		W
<i>Mnium hornum</i> Hedw.	F	W
<i>Mnium marginatum</i> (Dicks.) P.Beauv.	F	W
<i>Mnium stellare</i> Hedw.	F	W
<i>Neckera complanata</i> (Hedw.) Huebener	F	W
<i>Neckera crispa</i> Hedw.		W
<i>Neckera pumila</i> Hedw.		W
<i>Oligotrichum hercynicum</i> (Hedw.) Lam. & DC.	F	W
<i>Orthodontium lineare</i> Schwägr.	F	W
<i>Orthothecium intricatum</i> (Hartm.) Schimp.		W
<i>Orthothecium rufescens</i> (Dicks. ex Brid.) Schimp.		W
<i>Orthotrichum affine</i> Schrad. ex Brid.	F	W
<i>Orthotrichum anomalum</i> Hedw.	F	W
<i>Orthotrichum consimile</i> Mitt.		W
<i>Orthotrichum cupulatum</i> Hoffm. ex Brid.	F	W
<i>Orthotrichum diaphanum</i> Schrad. ex Brid.	F	W
<i>Orthotrichum ibericum</i> F.Lara & Mazimpaka	F	
<i>Orthotrichum lyellii</i> Hook. & Taylor	F	W
<i>Orthotrichum obtusifolium</i> Brid.	F	W
<i>Orthotrichum pallens</i> Bruch ex Brid.	F	W
<i>Orthotrichum patens</i> Bruch ex Brid.	F	W
<i>Orthotrichum pulchellum</i> Brunt.	F	W
<i>Orthotrichum pumilum</i> Sw. ex anon.	F	W
<i>Orthotrichum rivulare</i> Turner		W
<i>Orthotrichum rogeri</i> Brid.		W
<i>Orthotrichum rupestre</i> Schleich. ex Schwägr.	F	W
<i>Orthotrichum scanicum</i> Grönwall		W
<i>Orthotrichum schimperi</i> Hammar	F	W
<i>Orthotrichum shawii</i> Wilson		W
<i>Orthotrichum speciosum</i> Nees	F	W
<i>Orthotrichum sprucei</i> Mont.	F	W
<i>Orthotrichum stramineum</i> Hornsch. ex Brid.	F	W
<i>Orthotrichum striatum</i> Hedw.	F	W
<i>Orthotrichum tenellum</i> Bruch ex Brid.	F	W
<i>Oxyrrhynchium hians</i> (Hedw.) Loeske	F	W
<i>Oxyrrhynchium pumilum</i> (Wilson) Loeske	F	W
<i>Oxyrrhynchium schleicheri</i> (R.Hedw.) Röll	F	W
<i>Oxyrrhynchium speciosum</i> (Brid.) Warnst.	F	W
<i>Oxystegus tenuirostris</i> (Hook. & Taylor) A.J.E.Sm.		W
<i>Palustriella commutata</i> (Hedw.) Ochyra	F	W
<i>Palustriella falcata</i> (Hedw.) Hedenäs	F	W
<i>Paraleucobryum longifolium</i> (Hedw.) Loeske		W
<i>Phascum cuspidatum</i> Hedw.	F	W
<i>Philonotis arnellii</i> Husn.	F	W
<i>Philonotis caespitosa</i> Jur.	F	W
<i>Philonotis calcarea</i> (Bruch & Schimp.) Schimp.	F	W
<i>Philonotis fontana</i> (Hedw.) Brid.	F	W
<i>Philonotis marchica</i> (Hedw.) Brid.	F	W
<i>Philonotis seriata</i> Mitt.		W
<i>Physcomitrella patens</i> (Hedw.) Bruch & Schimp.	F	W
<i>Physcomitrium eurystomum</i> Sendtn.	F	W
<i>Physcomitrium pyriforme</i> (Hedw.) Bruch & Schimp.	F	W
<i>Physcomitrium sphaericum</i> (C.F.Ludw. ex Schkuhr) Brid.	F	B

<i>Plagiobryum zieri</i> (Hedw.) Lindb.		W
<i>Plagiomnium affine</i> (Blandow ex Funck) T.J.Kop.	F	W
<i>Plagiomnium cuspidatum</i> (Hedw.) T.J.Kop.	F	W
<i>Plagiomnium elatum</i> (Bruch & Schimp.) T.J.Kop.	F	W
<i>Plagiomnium ellipticum</i> (Brid.) T.J.Kop.	F	W
<i>Plagiomnium medium</i> (Bruch & Schimp.) T.J.Kop.	F	W
<i>Plagiomnium rostratum</i> (Schrad.) T.J.Kop.	F	W
<i>Plagiomnium undulatum</i> (Hedw.) T.J.Kop.	F	W
<i>Plagiopus oederianus</i> (Sw.) H.A.Crum & L.E.Anderson	F	W
<i>Plagiothecium cavifolium</i> (Brid.) Z.Iwats.	F	W
<i>Plagiothecium curvifolium</i> Schleph. ex Limpr.	F	W
<i>Plagiothecium denticulatum</i> (Hedw.) Schimp.	F	W
<i>Plagiothecium laetum</i> Schimp.	F	W
<i>Plagiothecium latebricola</i> Schimp.	F	W
<i>Plagiothecium nemorale</i> (Mitt.) A.Jaeger	F	W
<i>Plagiothecium succulentum</i> (Wilson) Lindb.	F	W
<i>Plagiothecium undulatum</i> (Hedw.) Schimp.	F	W
<i>Plasteurhynchium striatum</i> (Spruce) M.Fleisch.	F	W
<i>Platydictya jungermannioides</i> (Brid.) H.A.Crum		W
<i>Platygyrium repens</i> (Brid.) Schimp.	F	W
<i>Platypnnidium ripariooides</i> (Hedw.) Dixon	F	W
<i>Pleuridium acuminatum</i> Lindb.	F	W
<i>Pleuridium subulatum</i> (Hedw.) Rabenh.	F	W
<i>Pleurochaete squarrosa</i> (Brid.) Lindb.	F	W
<i>Pleurozium schreberi</i> (Willd. ex Brid.) Mitt.	F	W
<i>Pogonatum aloides</i> (Hedw.) P.Beauv.	F	W
<i>Pogonatum nanum</i> (Hedw.) P.Beauv.	F	W
<i>Pogonatum urnigerum</i> (Hedw.) P.Beauv.	F	W
<i>Pohlia andalusica</i> (Höhn.) Broth.		W
<i>Pohlia annotina</i> (Hedw.) Lindb.	F	W
<i>Pohlia bulbifera</i> (Warnst.) Warnst.	F	W
<i>Pohlia camptotrichela</i> (Renauld & Cardot) Broth.	F	W
<i>Pohlia cruda</i> (Hedw.) Lindb.		W
<i>Pohlia drummondii</i> (Müll.Hal.) A.L.Andrews		W
<i>Pohlia elongata</i> Hedw.		W
<i>Pohlia flexuosa</i> Hook.	F	W
<i>Pohlia lescuriana</i> (Sull.) Ochi	F	W
<i>Pohlia lutescens</i> (Limpr.) H.Lindb.	F	W
<i>Pohlia melanodon</i> (Brid.) A.J.Shaw	F	W
<i>Pohlia nutans</i> (Hedw.) Lindb.	F	W
<i>Pohlia wahlenbergii</i> (F.Weber & D.Mohr) A.L.Andrews	F	W
<i>Polytrichastrum alpinum</i> (Hedw.) G.L.Sm.		W
<i>Polytrichastrum formosum</i> (Hedw.) G.L.Sm.	F	W
<i>Polytrichastrum longisetum</i> (Sw. ex Brid.) G.L.Sm.	F	W
<i>Polytrichum commune</i> Hedw.	F	W
<i>Polytrichum juniperinum</i> Hedw.	F	W
<i>Polytrichum piliferum</i> Hedw.	F	W
<i>Polytrichum strictum</i> Menzies ex Brid.	F	W
<i>Polytrichum uliginosum</i> (Wallr.) Schrieb.	F	W
<i>Protobryum bryoides</i> (Dicks.) J.Guerra & M.J.Cano	F	W
<i>Pseudephemerum nitidum</i> (Hedw.) Loeske	F	W
<i>Pseudobryum cinclidoides</i> (Huebener) T.J.Kop.	F	W
<i>Pseudocalliergon lycopodioides</i> (Brid.) Hedenäs	F	
<i>Pseudocrossidium hornschuchianum</i> (Schultz) R.H.Zander	F	W
<i>Pseudocrossidium revolutum</i> (Brid.) R.H.Zander	F	W
<i>Pseudoleskeella catenulata</i> (Brid. ex Schrad.) Kindb.		W
<i>Pseudoleskeella nervosa</i> (Brid.) Nyholm		W

<i>Pseudoscleropodium purum</i> (Hedw.) M.Fleisch.	F	W	B
<i>Pseudotaxiphyllum elegans</i> (Brid.) Z.Iwats.	F	W	B
<i>Pterigynandrum filiforme</i> Hedw.	F	W	
<i>Pterogonium gracile</i> (Hedw.) Sm.		W	
<i>Pterygoneurum lamellatum</i> (Lindb.) Jur.		W	
<i>Pterygoneurum ovatum</i> (Hedw.) Dixon	F	W	
<i>Ptilium crista-castrensis</i> (Hedw.) De Not.	F	W	
<i>Ptychomitrium polyphyllum</i> (Dicks. ex Sw.) Bruch & Schimp.		W	
<i>Pylaisia polyantha</i> (Hedw.) Schimp.	F	W	B
<i>Racomitrium aciculare</i> (Hedw.) Brid.	F	W	
<i>Racomitrium affine</i> (F.Weber & D.Mohr) Lindb.		W	
<i>Racomitrium aquaticum</i> (Brid. ex Schrad.) Brid.		W	
<i>Racomitrium canescens</i> (Hedw.) Brid.	F	W	
<i>Racomitrium elongatum</i> Ehrh. ex Frisvoll	F	W	
<i>Racomitrium ericoides</i> (Brid.) Brid.	F	W	
<i>Racomitrium fasciculare</i> (Hedw.) Brid.		W	
<i>Racomitrium heterostichum</i> (Hedw.) Brid.	F	W	
<i>Racomitrium lanuginosum</i> (Hedw.) Brid.	F	W	
<i>Racomitrium obtusum</i> (Brid.) Brid.		W	
<i>Rhabdoweisia crenulata</i> (Mitt.) H.Jameson		W	
<i>Rhabdoweisia crispata</i> (Dicks.) Lindb.		W	
<i>Rhabdoweisia fugax</i> (Hedw.) Bruch & Schimp.		W	
<i>Rhizomnium pseudopunctatum</i> (Bruch & Schimp.) T.J.Kop.		W	
<i>Rhizomnium punctatum</i> (Hedw.) T.J.Kop.	F	W	B
<i>Rhodobryum ontariense</i> (Kindb.) Kindb.		W	
<i>Rhodobryum roseum</i> (Hedw.) Limpr.	F	W	
<i>Rhynchostegiella curviseta</i> (Brid.) Limpr.	F	W	B
<i>Rhynchostegiella tenella</i> (Dicks.) Limpr.	F	W	B
<i>Rhynchostegiella teneriffae</i> (Mont.) Dirkse & Bouman	F	W	
<i>Rhynchostegiella tenuicaulis</i> (Spruce) Kartt.		W	
<i>Rhynchostegium confertum</i> (Dicks.) Schimp.	F	W	B
<i>Rhynchostegium megapolitanum</i> (Blandow ex F.Weber & D.Mohr) Schimp.	F	W	
<i>Rhynchostegium murale</i> (Hedw.) Schimp.	F	W	B
<i>Rhynchostegium rotundifolium</i> (Scop. ex Brid.) Schimp.		W	
<i>Rhytidadelphus loreus</i> (Hedw.) Warnst.	F	W	
<i>Rhytidadelphus squarrosum</i> (Hedw.) Warnst.	F	W	B
<i>Rhytidadelphus subpinnatus</i> (Lindb.) T.J.Kop.		W	
<i>Rhytidadelphus triquetrus</i> (Hedw.) Warnst.	F	W	
<i>Rhytidium rugosum</i> (Hedw.) Kindb.	F	W	
<i>Sanionia uncinata</i> (Hedw.) Loeske	F	W	
<i>Schistidium apocarpum</i> (Hedw.) Bruch & Schimp.	F	W	B
<i>Schistidium confertum</i> (Funck) Bruch & Schimp.		W	
<i>Schistidium crassipilum</i> H.H.Bлом	F	W	
<i>Schistidium elegantulum</i> H.H.Bлом	F	W	
<i>Schistidium flaccidum</i> (De Not.) Ochyra		W	
<i>Schistidium helveticum</i> (Schkuhr) Deguchi		W	
<i>Schistidium papillosum</i> Culm.		W	
<i>Schistidium pruinosum</i> (Wilson ex Schimp.) G.Roth		W	
<i>Schistidium rivulare</i> (Brid.) Podp.	F	W	
<i>Schistidium trichodon</i> (Brid.) Poelt		W	
<i>Schistostega pennata</i> (Hedw.) F.Weber & D.Mohr.		W	
<i>Sciuro-hypnum flotowianum</i> (Sendtn.) Ignatov & Huttunen		W	
<i>Sciuro-hypnum oedipodium</i> (Mitt.) Ignatov & Huttunen	F		
<i>Sciuro-hypnum plumosum</i> (Hedw.) Ignatov & Huttunen	F	W	B
<i>Sciuro-hypnum populeum</i> (Hedw.) Ignatov & Huttunen	F	W	B
<i>Sciuro-hypnum reflexum</i> (Starke) Ignatov & Huttunen	F	W	
<i>Scleropodium cespitans</i> (Wilson ex Müll.Hal.) L.F.Koch.	F	W	

<i>Scleropodium touretii</i> (Brid.) L.F.Koch.		W
<i>Scopelophila cataractae</i> (Mitt.) Broth.	F	W
<i>Scorpidium cossonii</i> (Schimp.) Hedenäs	F	W
<i>Scorpidium revolvens</i> (Sw. ex anon.) Rubers	F	W
<i>Scorpidium scorpioides</i> (Hedw.) Limpr.	F	W
<i>Scorpiurium circinatum</i> (Bruch) M.Fleisch. & Loeske		W
<i>Seligeria acutifolia</i> Lindb.		W
<i>Seligeria calcarea</i> (Hedw.) Bruch & Schimp.	F	W
<i>Seligeria calycina</i> Mitt. ex Lindb.		W
<i>Seligeria donniana</i> (Sm.) Müll.Hal.		W
<i>Seligeria pusilla</i> (Hedw.) Bruch & Schimp.		W
<i>Seligeria recurvata</i> (Hedw.) Bruch & Schimp.		W
<i>Seligeria trifaria</i> (Brid.) Lindb.		W
<i>Sematophyllum demissum</i> (Wilson) Mitt.		W
<i>Sematophyllum substrumulosum</i> (Hampe) E.Britton	F	
<i>Sphagnum affine</i> Renaud & Cardot	F	W
<i>Sphagnum angustifolium</i> (C.E.O.Jensen ex Russow) C.E.O.Jensen	F	W
<i>Sphagnum auriculatum</i> Schimp.	F	W
<i>Sphagnum capillifolium</i> (Ehrh.) Hedw.	F	W
<i>Sphagnum centrale</i> C.E.O.Jensen	F	W
<i>Sphagnum compactum</i> Lam. & DC.	F	W
<i>Sphagnum contortum</i> Schultz	F	W
<i>Sphagnum cuspidatum</i> Ehrh. ex Hoffm.	F	W
<i>Sphagnum fallax</i> (H.Klinggr.) H.Klinggr.	F	W
<i>Sphagnum fimbriatum</i> Wilson	F	W
<i>Sphagnum flexuosum</i> Dozy & Molk.	F	W
<i>Sphagnum fuscum</i> (Schimp.) H.Klinggr.		W
<i>Sphagnum girgensohnii</i> Russow	F	W
<i>Sphagnum inundatum</i> Russow	F	W
<i>Sphagnum magellanicum</i> Brid.	F	W
<i>Sphagnum majus</i> (Russow) C.E.O.Jensen	F	W
<i>Sphagnum molle</i> Sull.	F	W
<i>Sphagnum obtusum</i> Warnst.	F	
<i>Sphagnum palustre</i> L.	F	W
<i>Sphagnum papillosum</i> Lindb.	F	W
<i>Sphagnum platyphyllum</i> (Lindb. ex Braithw.) Warnst.	F	
<i>Sphagnum quinquefarium</i> (Braithw.) Warnst.	F	W
<i>Sphagnum riparium</i> Ångstr.		W
<i>Sphagnum rubellum</i> Wilson	F	W
<i>Sphagnum russowii</i> Warnst.	F	W
<i>Sphagnum squarrosum</i> Crome	F	W
<i>Sphagnum subnitens</i> Russow & Warnst.	F	W
<i>Sphagnum subsecundum</i> Nees	F	W
<i>Sphagnum tenellum</i> (Brid.) Pers. ex Brid.	F	W
<i>Sphagnum teres</i> (Schimp.) Ångstr.	F	W
<i>Sphagnum warnstorffii</i> Russow		W
<i>Splachnum ampullaceum</i> Hedw.	F	W
<i>Splachnum sphaericum</i> Hedw.		W
<i>Straminergon stramineum</i> (Dicks. ex Brid.) Hedenäs	F	W
<i>Syntrichia calcicola</i> J.J.Amann	F	W
<i>Syntrichia laevipila</i> Brid.	F	W
<i>Syntrichia latifolia</i> (Bruch ex Hartm.) Huebener	F	W
<i>Syntrichia montana</i> Nees	F	W
<i>Syntrichia papillosa</i> (Wilson) Jur.	F	W
<i>Syntrichia princeps</i> (De Not.) Mitt.		W
<i>Syntrichia ruralis</i> (Hedw.) F.Weber & D.Mohr	F	W
<i>Syntrichia virescens</i> (De Not.) Ochyra	F	W

<i>Taxiphyllum wissgrillii</i> (Garov.) Wijk & Margad.		W	B
<i>Tayloria tenuis</i> (Dicks.) Schimp.		W	
<i>Tetraphis pellucida</i> Hedw.	F	W	B
<i>Tetraplodon minoides</i> (Hedw.) Bruch & Schimp.	F		
<i>Tetrodontium brownianum</i> (Dicks.) Schwägr.		W	
<i>Thamnobryum alopecurum</i> (Hedw.) Gangulee	F	W	B
<i>Thuidium assimile</i> (Mitt.) A.Jaeger		W	
<i>Thuidium delicatulum</i> (Hedw.) Schimp.	F	W	
<i>Thuidium recognitum</i> (Hedw.) Lindb.	F	W	
<i>Thuidium tamariscinum</i> (Hedw.) Schimp.	F	W	B
<i>Tomentypnum nitens</i> (Hedw.) Loeske	F	W	
<i>Tortella bambergeri</i> (Schimp.) Broth.		W	
<i>Tortella flavovirens</i> (Bruch) Broth.	F		
<i>Tortella inclinata</i> (R.Hedw.) Limpr.	F	W	
<i>Tortella inflexa</i> (Bruch) Broth.		W	
<i>Tortella nitida</i> (Lindb.) Broth.		W	
<i>Tortella tortuosa</i> (Hedw.) Limpr.	F	W	B
<i>Tortula atrovirens</i> (Sm.) Lindb.		W	
<i>Tortula canescens</i> Mont.	F	W	
<i>Tortula cernua</i> (Huebener) Lindb.	F		
<i>Tortula cuneifolia</i> (Dicks.) Turner	F		B
<i>Tortula inermis</i> (Brid.) Mont.		W	
<i>Tortula lanceola</i> R.H.Zander	F	W	
<i>Tortula marginata</i> (Bruch & Schimp.) Spruce	F	W	B
<i>Tortula modica</i> R.H.Zander	F	W	B
<i>Tortula mucronifolia</i> Schwägr.		W	
<i>Tortula muralis</i> Hedw.	F	W	B
<i>Tortula schimperi</i> M.J.Cano, O.Werner & J.Guerrra		W	
<i>Tortula subulata</i> Hedw.	F	W	B
<i>Tortula truncata</i> (Hedw.) Mitt.	F	W	B
<i>Tortula vahliana</i> (Schultz) Mont.	F	W	
<i>Trematodon ambiguus</i> (Hedw.) Hornsch.	F		
<i>Trichodon cylindricus</i> (Hedw.) Schimp.	F	W	B
<i>Trichostomum brachydontium</i> Bruch		W	
<i>Trichostomum crispulum</i> Bruch	F	W	B
<i>Uloa bruchii</i> Hornsch. ex Brid.	F	W	B
<i>Uloa coarctata</i> (P.Beauv.) Hammar	F	W	
<i>Uloa crispa</i> (Hedw.) Brid.	F	W	B
<i>Uloa hutchinsiae</i> (Sm.) Hammar		W	
<i>Uloa phyllantha</i> Brid.	F	W	
<i>Uloa rehmannii</i> Jur.		W	
<i>Warnstorffia exannulata</i> (Schimp.) Loeske	F	W	
<i>Warnstorffia fluitans</i> (Hedw.) Loeske	F	W	
<i>Warnstorffia pseudostraminea</i> (Müll.Hal.) Tuom. & T.J.Kop.	F		
<i>Warnstorffia sarmentosa</i> (Wahlenb.) Hedenäs		W	
<i>Weissia brachycarpa</i> (Nees & Hornsch.) Jur.	F	W	B
<i>Weissia condensa</i> (Voit) Lindb.		W	
<i>Weissia controversa</i> Hedw.	F	W	B
<i>Weissia longifolia</i> Mitt.	F	W	
<i>Weissia rostellata</i> (Brid.) Lindb.	F	W	
<i>Weissia rutilans</i> (Hedw.) Lindb.	F	W	
<i>Weissia squarrosa</i> (Nees & Hornsch.) Müll.Hal.	F	W	
<i>Zygodon conoideus</i> (Dicks.) Hook. & Taylor	F	W	B
<i>Zygodon forsteri</i> (Dicks.) Mitt.		W	
<i>Zygodon rupestris</i> Schimp. ex Lorentz	F	W	
<i>Zygodon stirtonii</i> Schimp. ex Stirt.	F	W	
<i>Zygodon viridissimus</i> (Dicks.) Brid.	F	W	B